

Abortion Incidence and Severity of Complications in Liberia

Key Findings from the 2021
National Study



April
2023



MINISTRY OF HEALTH
Republic of Liberia



African Population and
Health Research Center
Transforming lives in Africa through research

Table of Contents

List of Tables	4
List of Figures	4
Acknowledgments	5
Foreword	6
List of Acronyms	7
Executive Summary	9
Background	11
Data and Methods	13
Data Sources and Sampling Procedure	13
Data Analysis	15
Ethical Approval	16
Findings	17
Incidence of Induced Abortion	19
Abortion Rate and Ratios in Liberia	20
Unintended Pregnancy Rates and Induced Abortion	22
Severity of Abortion Complications	23
Discussion	25
Recommendations	26
References	27
Appendices	29

List of Tables

Table 1: Health facilities sampled in the Health Facility Survey, Liberia, 2021	17
Table 2: Estimated number of women treated for PAC in 2021 by regions in Liberia	17
Table 3: Unintended pregnancy and abortion	21
Table 4: Socio-demographic characteristics of patients	22
Table 5: Reproductive health characteristics of women seeking PAC, Liberia 2021	23

List of Figures

Figure 1: Regional distribution of PAC cases treated in health facilities	18
Figure 2: Estimated number of PAC patients in 2021 by facility level and ownership	18
Figure 3: Proportion of Induced Abortion Complications receiving treatment	19
Figure 4: Incidence of Induced Abortion, by region, Liberia, 2021	20
Figure 5: Abortion rate in Liberia (Number of induced abortion per 1,000 WRA)	21
Figure 6: Abortion ratio in Liberia (Number of induced abortion per 1,000 live births)	21
Figure 7: Severity of abortion complications among women treated in health facilities	24
Figure 8: Prevalence of abortion-related complications in health facilities	24

Acknowledgments

This report is based on research generously supported by the Swedish International Development Cooperation Agency (Sida) (Grant#. 12103) to the African Population and Health Research Center (APHRC). The study was led by the Ministry of Health, Liberia, in collaboration with the Clinton Health Access Initiative (CHAI Liberia), the APHRC and the Guttmacher Institute.

The study team wishes to thank the Ministry of Health, Liberia, particularly the Minister – Dr. Wilhelmina S. Jallah, the Deputy Minister/Chief Medical Officer- Dr. Francis N. Kateh, the Deputy Minister for Research, Planning and M&E- Hon. Varfley Tulay, the Director Family Health Program – Ms. Bentoe Z. Tehoungue, Director, Health Research – Nelson Dunbar for supporting the research. In addition, we are grateful to the Country Director of CHAI Liberia, Dr. Moses Massaquoi, for his leadership and support of this study. Further, we are indebted to all the staff at the Ministry of Health- Research unit for finding time to supervise data collection in the counties.

We are thankful to the heads of health facilities for the collaboration and support accorded during the study. Our appreciation also goes out to all the study participants: health providers, women and girls, and other professionals who agreed to participate in the research. We express gratitude to our highly professional fieldworkers for the dedication and excellence they demonstrated throughout the data collection period. Finally, we appreciate Martin Bangha, Issabelah Mutuku and Grace Kibunja for their technical support in the finalization of the report.



Foreword

In recent years, the commitments by the Government of Liberia to address the high maternal mortality in the country (through several interventions and guidelines that create an enabling environment for women to access high quality maternal health care) have yielded positive progress. For instance, according to the recent Liberia Demographic and Health Survey, maternal mortality ratio reduced from 1,072 deaths per 100,000 live births in 2013 to 742 deaths per 100,000 live births in 2019-2020. Even so, this progress has been slow with several existing gaps and weaknesses, especially in addressing abortion-related deaths.

While unsafe abortion is among leading causes of maternal illness and death, there is limited scientific information on the context in which women seek unsafe abortion, the magnitude of the problem and the resulting complications from such abortion.

This study, conducted by the Ministry of Health Liberia, the Clinton Health Access Initiative Liberia (CHAI), the African Population and Health Research Center (APHRC), and the Guttmacher Institute (GI) offers new and updated evidence on the incidence of induced abortion in Liberia, the profiles of women seeking post-abortion-related care in health facilities, the severity of abortion-related complications and the care they receive. The study provides timely information on a major, but preventable cause of death and suffering in our society today.

The findings in this report affirm that improving access to affordable and effective family planning and/or contraception is key to preventing unintended pregnancy and unsafe abortion. Effective family planning and/or contraception would also save the Government of Liberia the huge financial, personnel and other resources that currently go into the treatment of unsafe abortion and its associated complications. The report also drives home the importance of training to adequately equip our health providers with the requisite skills and knowledge to provide quality abortion-related care to women.

Taken together, this report is a wake-up call on the reality of unsafe abortion in Liberia and the need to urgently find a lasting solution to this preventable cause of maternal morbidity and mortality.

List of Acronyms

ACRE	Atlantic Center for Research and Evaluation
AICM	Abortion Incidence and Complications Methodology
APHRC	African Population and Health Research Center
CHAI	Clinton Health Access Initiative
FP	Family Planning
GFF	Global Financing Facility
HFS	Health Facility Survey
KIS	Knowledgeable Informant Survey
LISGIS	Liberia Institute of Statistics and Geo-Information Services
LDHS	Liberia Demographic and Health Survey
MOH	Ministry of Health
MPI	Multidimensional Poverty Index
PAC	Post-Abortion Care
PMS	Prospective Morbidity Survey
RMNCAH	Reproductive maternal, newborn, child and adolescent health
SGBV	Sexual and Gender-Based Violence
UL-PIRE	University of Liberia-Pacific Institute for Research and Evaluation
WHO	World Health Organization

Study Team

Ministry of Health, Liberia

Bentoe Z. Tehoungue, Principal Investigator
Nelson K. Dunbar, Co-Investigator

Clinton Health Access Initiative

Dr. Moses Massaquoi, Principal Investigator
Vekeh Donzo, Co-Investigator
Lily Lu, Co-Investigator
Matthew Nviiri, Co-Investigator

African Population and Health Research Center

Boniface Ushie, Principal Investigator
Kenneth Juma, Co-Investigator
Ramatou Ouedraogo, Co-Investigator
Esther Mutuku, Co-Investigator

Guttmacher Institute

Margaret Giorgio, Co-Investigator
Jesse Philbin, Co-Investigator
Stephanie Kung, Co-Investigator
Akin Bankole, Co-Investigator

Report Citation

Abortion incidence and severity of complications in Liberia: Key findings from the 2021 National Study (Monrovia, Liberia: Ministry of Health, Liberia, African Population and Health Research Center, Clinton Health Access Initiative Liberia, and Guttmacher Institute 2022).

Executive Summary


The World Health Organization (WHO) defines unsafe abortion as a procedure for terminating a pregnancy performed by persons lacking the necessary skills or an environment not in conformity with minimal medical standards, or both. Based on the current Liberian laws, abortion is only allowed under a limited set of conditions, including to save the life of a woman and to preserve physical or mental health, and when the pregnancy results from rape or incest or other felonious intercourse, and when the pregnancy has fetal impairment. Unsafe abortion remains a leading cause of maternal morbidity and death in Liberia while the treatment of resultant complications drains significant amounts of financial and material resources from the health system.

This report presents findings from a 2021 nationwide abortion study in Liberia, that aimed to estimate the incidence and severity of related complications. We conducted the study across all regions and counties in Liberia between September 2021 and March 2022, and in a nationally representative sample of public and private health facilities, including clinics, health centers and hospitals. The study used the abortion incidence complications methodology (AICM) to estimate abortion incidence and the severity of abortion complications in the country.

Findings showed that an estimated 38,779 induced abortion occurred in Liberia in 2021, translating to an induced abortion rate of 30.7 per 1,000 women of reproductive age, and an induced abortion ratio of 229 abortion per 1,000 live births. There were significant regional variations in the abortion rates, with North Central having the lowest abortion rate (6.6/1000) and South-Central having the highest (49.7/1000). In the same year, approximately 14,555 women received care for complications resulting from induced and spontaneous abortion in health facilities. The vast majority of women who sought post-abortion care services did so in public (67%) and primary-level health facilities (74%). The national unintended pregnancy rate was 86.5 per 1,000 WRA, with wide regional differences. The high abortion rates and unintended pregnancy rates in the South-Central region may imply that many women and girls have a high unmet need for family planning. According to the 2019-2020 Liberia Demographic and Health Survey, one-third of married women had an unmet need for family planning, 21% for spacing and 13% for limiting number of children. In line with expectations, the incidence rate is highest in regions with the highest unintended pregnancy rate further reinforcing the argument that legal restrictions do not reduce the occurrence of abortion, instead restrictions drive women toward unsafe abortion.

Findings also showed that women seeking abortion-related care in Liberian health facilities were socially and demographically diverse. Almost half of all patients seeking post-abortion care presented with either severe (37%) or near miss (11%) complications, and this varied widely by region. North-Western (52 %), North Central (43 %), and South-Central (40%) regions had the highest proportions of women with severe complications. South-Eastern B (17%) and South Eastern A (3%) regions had the lowest proportion of women with severe abortion-related complications, even though these two regions had the highest proportions of women with moderately severe complications. Most common complications reported among post-abortion patients were infections and sepsis (59%), hemorrhage (31%) and systolic blood pressure less than 91 mmHg (24%), and temperatures higher than 37.3°C (22%). There was one death during the study period.

Interventions to increase family planning while eliminating barriers to access especially for adolescents, reduce unmet need, and increasing use of safe abortion methods (e.g. using manual vacuum aspiration [MVA], misoprostol, or combipack) are effective approaches to reduce abortion-related morbidity and mortality. There is also great consensus that strengthening the ability of health facilities, especially the low-level health facilities to provide the basic post-abortion care services yields greater return on investment since the women who present at these facilities can get appropriate levels of quality care. Finally, there is need to reform the current abortion law to facilitate the reduction of unsafe abortion and also strengthen the fidelity of implementation of the clinical standards and guidelines for managing abortion-related complications within health facilities.



Background

The evolution of safe and effective medical procedures to perform safe abortion, along with providing universal access to such services, has the potential to eliminate unsafe abortion and related deaths entirely (1). However, unsafe abortion contributes about 13% of maternal death (2), with 25 million unsafe abortion occurring annually worldwide (3), majority of which are recorded in low and middle-income countries. Irrespective of the nature of abortion (spontaneous and or induced), and regardless of a country's abortion legal framework, post-abortion care (PAC) is endorsed globally as an essential emergency obstetric intervention to address complications associated with abortion (4,5).

Liberia, for a long time has had the highest maternal mortality ratios globally at 1,072 deaths per 100,000 live births in 2013 (6), and 742 deaths per 100,000 live births in the latest demographic and health survey (7). Most of these deaths are attributed to preventable and treatable complications, including hemorrhage (25%), hypertension (16%), unsafe abortion (10%), and sepsis (10%) (8). Deaths associated with unsafe abortion are often grossly underestimated, partly because of the social stigma that surrounds abortion and the fact that some deaths are not necessarily reported as linked to abortion (9). Under the Liberian penal code, abortion is only allowed under a limited set of conditions, including to save the life of the pregnant woman and to preserve her physical or mental health, when the pregnancy results from rape or incest or other felonious intercourse, and when the pregnancy has fetal impairment. However, two doctors must certify that a woman/girl meets these conditions before a pregnancy can be terminated, and only a doctor can perform the procedure.

Even with these restrictions, abortion, especially unsafe abortion remain common in Liberia, and previous abortion incidence estimates have ranged from 6% (Demographic and Health Survey) in 2007 (10) to 32% (population-based study) in 2015 (11). In Liberia, adolescent pregnancy rates remain high (30% adolescent pregnancy rate) (7), and is a contributor to unsafe abortion rates in the country, as unintended pregnancy is a precursor to induced abortion (12), and adolescents are particularly vulnerable to lack of safe abortion access (13).

There have been high-level commitments by the Government of Liberia and stakeholders to address the high rates of maternal deaths in the country. Part of these commitments are reflected in the inclusion of unsafe abortion-related maternal morbidity and mortality as a priority investment area under the Liberia Reproductive Maternal Newborn Child and Adolescent Health (RMNCAH) Investment Case of 2016-2020 (8). In December 2019, the Liberia MOH, with support from the CHAI, World Bank/GFF, and UNFPA, developed the first-ever National Comprehensive Abortion Care Guidelines that specify programmatic and clinical interventions for safe and legal abortion and PAC (14). Nevertheless, there still exist several gaps and weaknesses in addressing sexual and reproductive health (SRH) issues that contribute to abortion, such as limited access to modern and effective contraception and absence of comprehensive sexuality education. In 2019, stakeholders in Liberia (including lawmakers) began revisions to the country's Public Health Law (Title 33), that includes provisions on expanding access to safe abortion. As of July 2022, the revised Public Health Law had passed the Lower House of Parliament. If passed by the Upper House and assented into law, the Public Health Law will effectively repeal the 1976 Liberia Penal Code, thereby expanding access to safe abortion.

It is important to note that unsafe abortion is a significant contributor to maternal morbidity and mortality in Liberia (8). However, there is lack of comprehensive data on abortion and post-abortion care (PAC) in the country. This means that little is known about the incidence of abortion and abortion-related morbidity, as well as the number of women and girls who seek PAC. This information is crucial for effective planning, budgeting, and implementation of reproductive, maternal, newborn, child, and adolescent health (RMNCAH) intervention.

Moreover, ongoing legal reforms around the public health law in Liberia provide an opportunity to address unsafe abortion. Having robust and up-to-date evidence on the incidence of induced abortion, the severity of complications due to unsafe abortion, and the quality of abortion-related care in Liberia is necessary to consolidate gains made through the ongoing legal reforms. This information can also facilitate stronger campaigns and advocacy around the abortion discourse in Liberia and increase policy engagement to increase budgeting and investment in maternal health and post-abortion care.

This report presents the first national representative estimates of the incidence of induced abortion and the severity of abortion-related complications in Liberia. We have excluded data related to the quality of post-abortion care and the health system costs of providing post-abortion care from this report. This data will be presented later.

Data & Methods

Data Sources and Sampling Procedure

Data used to estimate induced abortion incidence, the magnitude, and the severity of abortion complications was generated from three (3) surveys that make up the Abortion Incidence and Complications Methodology (AICM). These are the Health Facilities Survey (HFS), Knowledgeable Informant Survey (KIS), and Prospective Morbidity Survey (PMS). All fieldwork was conducted between September 2021 and March 2022.

Health Facility Survey (HFS)

The HFS is a national representative, stratified and randomly sampled survey of health facilities capable of providing PAC services as per the current National Guidelines for Comprehensive Abortion Care (2019). Thus, the sample frame included all clinics, health centers, and hospitals. Stratification was done by the five (5) geographical regions (South Central, North Western, North-Central, South Eastern A and South Eastern B), then by level of health facility (clinic, health center, hospital). These sampled public facilities were supplemented by purposely sampled private facilities (including private-for-profit, faith-based, and concession facilities) across the five (5) regions. The MOH Master Facility List obtained May 31, 2021, was used to identify eligible facilities from both the public and private sector. The universe included 894 facilities comprising clinics (802), health centers (59), and hospitals (34). We sampled a total of 132 health facilities for the HFS component of study, that included 31 hospitals (91.2%), 41 health centers (69.4%) and 60 clinics (7.5%). One hundred and twenty-eight (128) (97%) of the sampled facilities completed the HFS. We excluded three (3) hospitals because one burnt down during the study while two (2) were specialized hospitals.

The HFS interviews were conducted with a health professional who is knowledgeable about post-abortion care at each selected facility. In large facilities, such as hospitals, the HFS was completed by the head of the maternal health or labor and delivery unit or an obstetrician/gynecologist who oversees PAC services. In lower-level facilities, we administered the HFS to a nurse, midwife, officer in charge (OIC), or health worker who was able to provide information about abortion-related care in that facility. Each respondent was interviewed using a face-to-face structured questionnaire.

The HFS collected retrospective estimates of the number of in- and out-patient PAC cases treated at each facility in the past month and typical month, which helped in generating facility caseloads. In the event the respondent was unable to give the monthly estimates, retrospective estimates of the past year and typical year were requested. We used the average or typical month (year) to iron out the seasonality in abortion. Other information captured in the survey included the health facilities' provision of PAC services, including PAC management approaches, provider knowledge on CAC, and PAC family planning.

Knowledgeable Informants Survey (KIS)

The KIS involved in-person structured interviews with a sample of purposely selected respondents with knowledge about the provision of abortion and PAC in the country (nationally or pertaining to specific regions). Informants were not restricted to clinicians providing direct CAC services to patients. Potential key informants were identified through purposeful and snowball sampling.

The initial list included individuals who had one or more of the following roles:

1. Key MOH/County Health Team staff involved with SRH programs;
2. Head/informed member(s) of partner organizations implementing SRH programs;
3. Head/informed member(s) of SRH advocacy groups;
4. Researchers of sexual and reproductive health and rights (SRHR) from different regions of Liberia. The selected individual were asked to list up to two (2) referrals they believed were knowledgeable on the provision of abortion and PAC in Liberia.

A total of 89 key informants were interviewed on topics ranging from their perception regarding the type of providers women seek abortion from, the likelihood women will experience complications that require treatment in a facility according to the type of abortion provider used, to the likelihood that women who need treatment will receive it at a health facility. These questions were asked for four (4) major sub-groups of women within the population: rural poor, rural non-poor, urban poor, and urban non-poor. This information was used in the calculation of the multiplier. The multiplier is the number of women/girls who have induced abortion who either did not have complications, or had complications that were not treated in the formal health system, for everyone that received facility-based care.

Other data sources used for calculating abortion incidence were the 2019/2020 Liberia Demographic and Health Survey (LDHS) (7), which provided information on fertility, contraceptive prevalence, unmet need for contraception, birth wantedness, and measures of access to health care. We also drew data from 2021 population projection for the number of women aged 15-49 in the different regions of Liberia. We used the Poverty & Equity Brief for Liberia 2021(15) and the Global Multidimensional Poverty Index (MPI) Country Briefing 2021: (16), to estimate the proportion of poor/non-poor in urban/rural settings in Liberia.

Prospective Morbidity Survey (PMS)

The PMS is a nationally representative survey that targeted a sub-sample of the HFS sample. One hundred (100) health facilities were selected from the HFS sample of 128 to participate in the PMS.

The PMS captured data on each woman who obtained post-abortion care (and their provider) at the selected health facilities during a 30-day period, for the women who consented. The inclusion criteria were all women seeking treatment for post-abortion complications and whose pregnancy termination occurred at 28 weeks or lower gestational age. However, we excluded all women who presented with ectopic and molar pregnancies, blighted ovum, and threatened abortion. These cases were excluded because the clinical management of the first three (3) is evacuation (we did not want to confuse that with abortion) and the last because a woman with threatened abortion may still leave the facility with her pregnancy continuing.

Trained clinicians and a few non-clinicians were trained and stationed at each selected facility for the 30 days of observation to collect data as women prospectively presented and were treated for PAC. In high-volume facilities, more than one enumerator was trained to collect data.

The PMS collected information on PAC patients' socio-demographic characteristics, reproductive and clinical histories, diagnosis, clinical procedures, and other services performed to treat the woman at the facility, post-abortion contraception provision, and clinical management outcomes. Data from the PMS is specifically applied to the classification of the severity of abortion complications. Data was collected from 401 PAC patients and 432 PAC providers across these 100 health facilities.

Data Analysis

Incidence of Induced Abortion

To estimate the incidence of induced abortion, we used the HFS to first obtain the number of women treated for abortion-related complications in a year in each facility by averaging the yearly estimates of the number treated in the past month and an average month for in-patients and out-patients and adding the two estimates. These facility-level estimates were then weighted to generate national representative data, accounting for sampling proportions and non-response. Table 2 shows the annual caseloads by facility level and region, after which we excluded the referrals to avoid double counting PAC cases. Next, we adjusted the PAC case totals to exclude cases of spontaneous abortion (i.e. miscarriages) at the regional level, using the assumption that 3.41% of live births in each of the five (5) geographic regions are due to late-term miscarriages (at 13-24 weeks) (17), which are the ones that are likely to require facility-based treatment, to estimate the number of miscarriages per region (Appendix 1). We then subtracted that number (miscarriages) from the total number of PAC cases to obtain the number of PAC cases due to induced abortion. Finally, the weighted regional estimate of the number of women treated for induced abortion complications are added to generate the total number of women treated in health facilities for complications resulting from induced abortion in Liberia in 2021.

Responses from the KIS were used to calculate the “multiplier”, which represents the estimated proportion of all women who had an induced abortion that received treatment for abortion-related complications, including considerations that some women may not have complications, or may not seek care in the formal health system. The multiplier or adjustment factor is obtained as the inverse of that weighted proportion who seek care. The multiplier indicates, for every woman who had induced abortion complications and received care from a health facility, the number of additional women who had induced abortion who did not have complications or had complications but did not seek PAC in a health facility. Applying the multiplier to the facility PAC cases due to induced abortion provides the total estimated number of women who obtained an induced abortion in Liberia in 2021. From this number, the abortion incidence (number of induced abortion per year), abortion rate (number of induced abortion per 1,000 women of reproductive age [15-49]), and abortion ratio (number of induced abortion per 1,000 live births).

Severity of Abortion Complications

To calculate the severity of abortion complications, we used data from the PMS as well as the categorization of complications based on prior studies in the region on abortion-related complications (Box 1) (18-21). Each patient’s clinical signs and symptoms, diagnosis, and interventions are examined to generate five (5) distinct categories of complications – mild, moderately severe, severe, near-miss, and death, and this is presented by regions.

Box 1: <i>Criteria for classification of abortion-related morbidity</i>
Mild morbidity (requires all criteria)
Temperature 35.1°C-38.9°C with no clinical signs of infection* No system or organ failure Systolic blood pressure ≥90mm Hg Hemorrhage not requiring any transfusion
Moderately severe morbidity (requires ≥1 criterion)
Temperature 37.3°C-38.9°C Clinical signs of infections No organ or system failure No signs of shock Hemorrhage not requiring any transfusion
Severe morbidity (requires ≥1 criterion)
Temperature ≥39°C or ≤35°C and a clinical sign of infection Sepsis/septicemia with no signs of septic shock Pelvic abscess or pelvic peritonitis with no sign of shock Clinical anemia without hemorrhagic shock Uterine perforation without laparotomy, repair of uterine perforation, gut perforation, hysterectomy
Near-miss (requires ≥1 criterion)
Hemorrhagic shock Septic shock Generalized peritonitis Uterine perforation with laparotomy, repair of uterine perforation, gut perforation, or hysterectomy Organ/system failure Massive blood transfusion
Death
Loss of the life of a woman because of an abortion complication



Ethical Approval

This study was approved by the University of Liberia-Pacific Institute for Research and Evaluation Institutional Review Board (UL-PIRE) (now the Atlantic Center for Research and Evaluation (ACRE) Institutional Review Board, Protocol #21-07-275. In line with CHAI's internal policy, the study protocol and materials were also approved by the Clinton Health Access Initiative's internal Scientific and Ethical Review Committee (SERC). The APHRC Institutional Review Board also reviewed and approved the study. All investigators who worked on the team completed the human subjects' protection training before engaging in the study. All respondents provided signed (HFS and KIS) or verbal (PMS) informed consent before participation.

Findings

PAC Caseload

Table 1 presents the universe of health facilities with the potential to provide PAC in Liberia, the health facilities sampled, those that declined, and the total health facilities that participated in the HFS. Of the 894 unweighted universe of health facilities, 132 were sampled and 128 participated in the HFS. All sampled hospitals participated (100%), while health centers had a response rate of 97.6% and clinics (95.0%).

Table 1: Health facilities sampled in the Health Facility Survey, Liberia, 2021

Facility Type	Universe of health facilities with potential capacity to provide PAC	Health facilities sampled	Health facilities that declined participation	Health Facilities participating
Hospital	34	31	0	31 (100%)
Health center	59	41	1	40 (97.6%)
Clinic	802	60	3	57 (95.0%)
Total	894	132	4	128 (97.0%)

Findings from the HFS showed that 14,555 women received post-abortion care, including for induced and spontaneous abortion within health facilities in Liberia in 2021 (Table 2). Across regions, South Central (Montserrado, Margibi and Grand Bassa) had the highest number of cases (7,462), while the least PAC cases were seen in Southeast B (702). After excluding referrals and miscarriages (spontaneous abortion) from the total PAC caseloads, we estimated that the total number of women treated for induced abortion complications at health facilities was 8,461 (Table 2). Figure 1 shows the regional distribution of PAC cases due to induced abortion.

Table 2: Estimated number of women treated for PAC in 2021 by regions in Liberia

	Total # PAC cases (a)	# referrals (b)	# miscarriages (c)	# PAC cases due to induced abortion (a-b-c)
National	14,555	332	5,762	8,461
North Central	3,593	149	2,287	1,157
North Western	1,885	50	581	1,254
South Central	7,462	115	2,104	5,243
Southeast A	913	15	442	456
Southeast B	702	3	348	351




Figure 1: Regional distribution of PAC cases treated in health facilities

Most women presenting for PAC (74% or 10,767 cases) were treated at lower-level facilities (clinics), followed by hospitals (13.9% or 2,025 cases) (Figure 2). In addition, most women were treated in public health facilities (67% or 9,752 cases) compared to 33% or 4,803 cases in private facilities, highlighting that it is the public facilities that are common access points for patients seeking PAC services in Liberia.




Figure 2: Estimated proportion of women treated for PAC in 2021 by facility level and ownership

38,779
incidence
of induced
abortion in
Liberia in
2021. This
translates to
an abortion
rate for Liberia
at

30.7
per


1,000
women of
reproductive
age.



Post-abortion Care-seeking

Figure 3 shows the proportion of women with complications from induced abortion in Liberia who received post-abortion care in health facilities in 2021 by region of care. Just under a quarter of women (22%) at the national level with induced abortion complications received care in a facility.

Across the regions, North Central region (Bong, Lofa, Nimba counties) had the highest proportion of induced abortion complications treated in a facility (41%), and this could possibly be linked to more abortion induced through 'traditional' methods (known to be unsafe) within rural settings. At the same time, the South-Central region (Montserrado, Margibi, GB) had only 18% of induced abortion complications treated in a facility. This low proportion of women with induced abortion complications seeking care in South Central could reflect the large proportion of women in urban settings who have no complications from induced abortion (i.e. because safe abortion methods are more readily accessible).



Incidence of Induced Abortion

The national multiplier was 4.5, indicating that approximately 22% of all women who had induced abortion in Liberia experienced complications and sought treatment at a health facility.

Once we applied the multiplier to the PAC caseload numbers, we estimated that the incidence of induced abortion was 38,779 in Liberia in 2021. This translates to an abortion rate for Liberia at 30.7 per 1,000 women of reproductive age.

Figure 4 presents the regional estimates of the incidences of induced abortion in Liberia in 2021.

There exist substantial regional variations in the abortion rate, with the highest rate seen in the South Central region (49.7 per 1,000 women in reproductive age), followed by

North Western (42.5 per 1,000 women in reproductive age); and the lowest incidence was seen in North Central (6.6 per 1,000 women in reproductive age). However, we note that the regional incidence must be cautiously interpreted because it does not necessarily mean that the people who had these induced abortion are domicile in South Central; for example, it is possible that South Central may see more induced abortion due to accessible, safe abortion services, whether performed formally or informally.

Induced abortion by regions

11%

North Western

7%

North Central

74%

South Central

4%

Southeast A

3%

Southeast B





Figure 4: Incidence of induced abortion by regions, Liberia, 2021

Abortion Rate and Ratios in Liberia

Figure 5 shows the abortion incidence rates. Liberia's abortion rate is 30.7 abortion per 1,000 women of reproductive age (WRA). There are substantial regional variations in the abortion incidence. The abortion rates were highest in the South Central (49.7/1,000), perhaps due to accessible safe abortion services, whether performed formally or informally. North Western also recorded a high abortion rate (42.5/1,000) possibly because of the low regional family planning (FP) use, a high unmet need for FP, a high unintended pregnancy rate, and service inaccessibility.

Figure 6 shows the abortion ratios in Liberia – that is, the number of induced abortion per 1,000 live births. Liberia's abortion ratio is 229 abortion per 1,000 live births and there are significant regional variations in the abortion ratios.




Figure 5: Abortion rate in Liberia (Number of induced abortion per 1,000 WRA)




Figure 6: Abortion ratio in Liberia (Number of induced abortion per 1,000 live births)

¹ Number of pregnancies per 1,000 WRA

² Number of unintended pregnancies per 1,000 WRA, with unintended pregnancies defined as unintended births plus abortion

Unintended Pregnancy Rates and Induced Abortion

Table 3 presents the comparison between pregnancy rates, unintended pregnancy rates, proportion of pregnancies that are unintended, and the proportion of unintended pregnancies ending in abortion.

Of all pregnancies in Liberia in 2021, 52% were unintended, and of all these unintended pregnancies, 35% ended in abortion (Table 3). The national unintended pregnancy rate was 86.5 per 1,000 WRA. The highest unintended pregnancy rates were in North-Western (118.0 per 1,000 WRA) and South Central (99.8 per 1,000 WRA) regions. Similarly, South Central (Montserrado, Margibi, GB) had the highest proportion of pregnancies that are unintended, compared to the national estimate (64% vs. 52%). South Central also had the highest proportion of unintended pregnancies ending in abortion, above the national estimate (50% vs. 35%). *Appendix 2 details the outcomes of all pregnancies in Liberia in 2021.*

Table 3: Unintended pregnancy and abortion

	Pregnancy rate	Unintended pregnancy rate	Proportion of pregnancies that are unintended	Proportion of unintended pregnancies ending in abortion
National	194.7	86.5	52%	35%
North Central	195.6	66.0	40%	10%
North Western	244.9	118.0	57%	36%
South Central	183.3	99.8	64%	50%
Southeast A	215.5	78.3	43%	25%
Southeast B	186.5	65.3	42%	26%

Characteristics of PAC Patients

A total of 401 women were interviewed during the 30-day PMS data collection. Table 4 presents the socio-demographic characteristics of the respondents. Half (50%) of all PAC patients were young women aged (< 25 years), while about 10% were 35 years or older. Most women were from urban settings (63%), married or in a union (67%), were working (employed/self-employed) (56%), and had primary-level education (39%).

Table 4: Socio-demographic characteristics of PAC patients

Patient's Background History (n=401)		%
Age	13-19	23%
	20-24	27%
	25-34	40%
	35 and above	10%
Residence	Urban	63%
	Rural	37%
Marriage	In union	67%
	Not in union	32%
	Refused to disclose	1%
Employment/Work	Working, employed or self-employed	56%
	Unemployed	35%
	Other (housewife; student)	9%
Highest level of education	No education	20%
	Primary	39%
	Secondary	27%
	Tertiary	14%

*Weighted percentages

Table 5 presents the reproductive health characteristics of women seeking PAC in Liberian health facilities. Most women reported one or two total pregnancies (44%). Among 275 women with previous deliveries, 56% reported having had between 1-2 total live births. Among women with biological living children, the majority had one or two biological living children (54%). Further, of the 119 women with previous experience of spontaneous abortion, 61% reported once, while 39% had at least two episodes of spontaneous abortion. About 36% of the women in the study did not want the current pregnancy for which they were seeking PAC as at the time they became pregnant, and a majority (53%) were not doing something to prevent or delay the pregnancies.

Table 5: Reproductive health characteristics of women seeking PAC, Liberia 2021


Reproductive health history (n=401)	Response options	%
Total pregnancies (n=401)	1-2	44%
	3-4	35%
	5-6	12%
	7 and above	10%
Total number of children given birth to previously (n=275)	1-2	56%
	3-4	28%
	5 and above	17%
Total biological living children (n=271)	None	4%
	1-2	54%
	3-4	27%
	5 and above	14%
Previous spontaneous abortion (n=119)	1	61%
	2	28%
	3 and above	11%
Previous induced abortion (n=67)	1	51%
	2	41%
	3 and above	8%
Pregnancy Wantedness (n=401)	Wanted the pregnancy and to carry it to term	60%
	Wanted but changed mind	4%
	Did not want it at all	30%
	Had no choice	6%
Doing something to delay/prevent pregnancy (n=401)	No	53%
	Yes	47%

*Weighted percentages

Severity of Abortion Complications


Based on the categorization of complications (see Box 1) for women who presented for PAC during the 30 days of facility observation, about 26% of women seeking PAC presented with mild complications, 25% with moderately severe, and 37% with severe complications (Figure 7). Nationally, 11% of all complications were near-miss or resulted in death.

Across the regions, North-Western (52%), North Central (43%), and South-Central (40%) regions had the highest proportions of women with severe complications. South Eastern B (17%) and South Eastern A (3%) regions had the least proportion of women with severe abortion-related complications, even though these two regions had the highest proportion of women with moderately severe complications. The regional variations in the severity of complications shown in Figure 7 could be reflective of differences in facility infrastructure, provider capacity and knowledge of clinical practice, availability of supplies and commodities.



Abortion-Related Complications

Figure 8 shows the distribution of specific abortion-related complications by women presenting for PAC services. More than half of women were diagnosed with both clinical signs of infections and sepsis (59%), about one-third had hemorrhage (31%) and slightly over a fifth of the women had a systolic blood pressure less than 91 mmHg (24%), and temperatures higher than 37.3°C (22%). There was one death during the 30-day study observation period.



Discussion

The AICM approach suggests that approximately 38,779 abortion occurred in Liberia in 2021, corresponding to a national abortion rate of 30.7 per 1,000 WRA. The estimated abortion rate is largely comparable to existing abortion rate estimates (22). For instance, 2017 modeling from the Guttmacher Institute found that the abortion rate in western Africa was 31 per 1,000 WRA between 2010-2014 (23). Studies in neighboring countries have estimated the induced abortion incidence rate to be 25 per 1,000 in Burkina Faso (24), 33 per 1,000 in Nigeria (25), 44 per 1,000 in Ghana (26), and 16 per 1000 in Senegal (27). It is however valuable to highlight that these previous studies were conducted close to a decade ago.

There are significant variations in the abortion rate across regions in Liberia, with North Central having the lowest abortion rates (6.6/1,000) and South-Central having the highest (49.7/1,000). The South-Central region also reported a high unintended pregnancy rate (99.8/1,000). The high abortion rates and unintended pregnancy rates in the South-Central region may indicate that women and girls have high unmet need for family planning. For instance, in 2020, one-third of married women had an unmet need for family planning, 21% for spacing and 13% for limiting number of children, and this gap increases among adolescents and young people (7). Further, the two (2) regions (North-Western and South-Central) with the highest unintended pregnancy rates also had the highest abortion incidence rates. This reinforces the argument that legal restrictions do not necessarily reduce the occurrence of abortion, but rather drive women toward unsafe abortion methods likely to result in complications and death (22). Interventions to increase family planning, reduce unmet need, and to increase use of safe abortion methods (e.g. using manual vacuum aspiration [MVA], misoprostol, or combipack) are effective approaches to reduce abortion-related morbidity and mortality (28).

Data from the HFS indicates that the vast majority of women who seek PAC services do so in public (67%) and primary-level health facilities (74%) compared to health centers and hospitals. Yet, despite these public and primary-level facilities being the dominant access points to women seeking PAC services in Liberia, they are most often the least equipped with trained staff, essential equipment, commodities, and supplies for PAC, thus impeding access to quality and comprehensive abortion-related services (29).

Study findings show that women seeking abortion-related care in Liberian health facilities were socially and demographically diverse. This study shows that almost half of the patients seeking post-abortion care presented with either severe (37%) or near-miss (11%) complications and with wide regional variations. The PAC cases with this level of severity were higher than those documented in some studies, like Zimbabwe and Uganda (19-20%) (30,31) but comparable to the level 37% found in Kenya and Sierra Leone (18). Safe abortion, i.e., procedure for terminating an unintended pregnancy carried out by providers with the necessary knowledge, skills, and qualifications in an environment that conforms to minimal medical standards (e.g. following WHO guidelines) and with appropriate commodities, equipment and supplies, typically result in extremely low rates of complications. Unsafe abortion however, results in severe complications, some of which are life-threatening. Women with severe complications often require admission into intensive and high dependency care units, long admission periods, and treatment by highly skilled health providers, such as gynecologists who are often scarce in low-income countries, especially at lower facility levels (31). Due to the associated morbidity and mortality, the burden of unsafe abortion on the health system are enormous (32).

While there are ongoing legislative processes in Liberia to reform the existing laws around abortion and enhance access to safe abortion care and reduce unsafe abortion and improve maternal health in general, there remains a need to implement and routinely monitor the use of existing clinical guidance for delivering PAC and addressing abortion-related complications in Liberia.

Recommendations

- 01 Accelerate the legal and policy reforms that address the causes of unsafe abortion and unintended pregnancies and eliminate barriers to access of sexual and reproductive health services for all women and girls. Liberia has already committed to domesticating the Maputo Protocol, that creates a framework for addressing sexual and gender-based violence (SGBV), facilitating a return to school criteria for pregnant and parenting adolescents, safe abortion care and the age of consent for sex and marriage, among others.
- 02 Access to quality family planning services and effective modern methods for all women and men.
- 03 Expanding access to quality post-abortion care, including post-abortion contraception counseling and method provision at all health system levels. This also includes strengthening the capacity of lower-level health facilities and mid-level providers to utilize appropriate uterine evacuation technologies, including MVA, misoprostol, and mifepristone/misoprostol combipack, availing essential equipment and supplies, and ensuring pre-service and in-service training of providers on PAC. The 2019 National CAC Guidelines should be disseminated and used to support training and mentoring of all providers on high-quality PAC.
- 04 Ensure full dissemination and implementation of the revised Public Health Law once passed to all communities, health facilities, and policymakers.
- 05 Community education and awareness on the burdens and dangers of unsafe abortion, current legal provisions on abortion and current access points (e.g. hospitals), stigma reduction, and awareness on a wide range of family planning methods available in Liberia.
- 06 Strengthen the health system and health facilities to ensure that they have the necessary equipment and supplies, including medication and contraception, to provide quality sexual and reproductive health services, including abortion and post-abortion care.
- 07 Collaborate with local and international organizations, including NGOs, to support and implement programs that promote access to comprehensive sexual and reproductive health services, including family planning, abortion care, and post-abortion care.


References

1. World Health Organization. Safe abortion: Technical & policy guidance for health systems Legal and policy considerations. Geneva; 2012
2. World Health Organization. Unsafe abortion: Global and regional estimates of the incidence of unsafe abortion and associated mortality in 2008. Geneva; 2009.
3. Guttmacher Institute. Induced Abortion Worldwide. 2018. 2018 Mar
4. Temmerman M. Missed opportunities in women's health: Post-abortion care. Lancet Glob Health. 2019 Jan 1;7(1):e12-3.
5. DeJong J. The role and limitations of the Cairo International Conference on Population and Development. Soc Sci Med .2000;51(6):941-53.
6. Liberia Institute of Statistics and Geo-Information Services (LISGIS) Monrovia L, Ministry of Health and Social Welfare Monrovia L, National AIDS Control Program Monrovia L, ICF International Inc. Rockville MU. Liberia Demographic and Health Survey 2013. Monrovia; 2014 Aug.
7. Liberia Institute of Statistics and Geo-Information Services (LISGIS), Ministry of Health, The DHS Program ICF. Liberia Demographic and Health Survey 2019-20. Monrovia; 2021 Aug.
8. Ministry of Health L. Investment Case for Reproductive, Maternal, New-Born, Child, and Adolescent Health 2016-2020. Monrovia; 2016.
9. Rasch V. Unsafe abortion and postabortion care - an overview. Acta Obstet Gynecol Scand [Internet]. 2011 Jul 1;90(7):692-700.
10. Liberia Institute of Statistics and Geo-Information Services (LISGIS), Ministry of Health and Social Welfare, National AIDS Control Program, Macro International Inc. Calverton MU. Liberia Demographic and Health Survey 2007. Monrovia; 2008 Jun.
11. Moseson H, Massaquoi M, Dehlendorf C, Bawo L, Dahn B, Zolia Y, et al. Reducing under-reporting of stigmatized health events using the List Experiment: Results from a randomized, population-based study of abortion in Liberia. Int J Epidemiol. 2015 Dec 1;44(6):1951-8.
12. Atuhaire S. Abortion among adolescents in Africa: A review of practices, consequences, and control strategies. Int J Health Plann Manage. 2019 Oct 1;34(4):e1378-86.
13. World Health Organization. Adolescent pregnancy. Fact sheets. 2022.
14. Liberia Ministry of Health, Clinton Health Access Initiative L. National Guidelines for Comprehensive Abortion Care: Safe abortion for legal indications & post-abortion care. Monrovia; 2019.
15. World Bank Group. Poverty & Equity Brief Liberia. 2021 Apr.
16. Oxford Poverty and Human Development Initiative (OPHI). Global MPI Country Briefing 2022: Liberia (Sub-Saharan Africa). Oxford; 2022 Oct.

17. Harlap S, Shiono PH, Ramcharan S. Spontaneous foetal losses in women using different contraceptives around the time of conception. *Int J Epidemiol*. 1980 Mar;9(1):49-56.
18. Ministry of Health Kenya, African Population and Health Research Center (APHRC), Guttmacher Institute. Incidence and Complications of Unsafe Abortion in Kenya: Key findings of a national study. Nairobi; 2013.
19. Liyew EF, Yalew AW, Afework MF, Essén B. Incidence and causes of maternal near-miss in selected hospitals of Addis Ababa, Ethiopia. *PLoS One*. 2017 Jun 6;12(6):e0179013.
20. Owolabi OO, Cresswell JA, Vwalika B, Osrin D, Filippi V. Incidence of abortion-related near-miss complications in Zambia: Cross-sectional study in Central, Copperbelt and Lusaka Provinces. *Contraception* [Internet]. 2017 Feb 1;95(2):167-74.
21. Owolabi O, Riley T, Juma K, Mutua M, Pleasure ZH, Amo-Adjei J, et al. Incidence of maternal near-miss in Kenya in 2018: Findings from a nationally representative cross-sectional study in 54 referral hospitals. *Sci Rep*. 2020 Dec 16;10(1):15181.
22. Bearak J, Popinchalk A, Ganatra B, Moller AB, Tunçalp Ö, Beavin C, et al. Unintended pregnancy and abortion by income, region, and the legal status of abortion: Estimates from a comprehensive model for 1990-2019. *Lancet Glob Health*. 2020 Sep 1;8(9):e1152-61.
23. Bearak JM, Popinchalk A, Beavin C, Ganatra B, Moller AB, Tunçalp Ö, et al. Country-specific estimates of unintended pregnancy and abortion incidence: A global comparative analysis of levels in 2015-2019. *BMJ Glob Health*. 2022 Mar 1;7(3):e007151.
24. Sedgh G, Rossier C, Kaboré I, Bankole A, Mikulich M. Estimating Abortion Incidence in Burkina Faso Using Two Methodologies. *Stud Fam Plann*. 2011 Sep 1;42(3):147-54.
25. Bankole A, Adewole IF, Hussain R, Awolude O, Singh S, Akinyemi JO. The Incidence of Abortion in Nigeria. *Int Perspect Sex Reprod Health*. 2015 Dec;41(4):170-81.
26. Keogh SC, Otupiri E, Chiu DW, Polis CB, Hussain R, Bell SO, et al. Estimating the incidence of abortion: a comparison of five approaches in Ghana. *BMJ Glob Health*. 2020 Apr 1;5(4):e002129.
27. Sedgh G, Sylla A, Philbin J, Keogh S, Ndiaye S. Estimates of the Incidence of Induced Abortion And Consequences of Unsafe Abortion in Senegal. *Int Perspect Sex Reprod Health*. 2015 Mar 1;41:11-9.
28. Ministry of Health Z. Standards and guidelines for reducing unsafe abortion morbidity and mortality in Zambia. 2009.
29. Owolabi OO, Biddlecom A, Whitehead HS. Health systems' capacity to provide post-abortion care: a multicountry analysis using signal functions. *Lancet Glob Health*. 2019 Jan 1;7(1):e110-8.
30. Madziyire MG, Polis CB, Riley T, Sully EA, Owolabi O, Chipato T. Severity and management of postabortion complications among women in Zimbabwe, 2016: A cross-sectional study. *BMJ Open*. 2018 Feb 1;8(2):e019658.
31. Qureshi Z, Mehrtash H, Kouanda S, Griffin S, Filippi V, Govule P, et al. Understanding abortion-related complications in health facilities: Results from WHO multicountry survey on abortion (MCS-A) across 11 sub-Saharan African countries. *BMJ Glob Health*. 2021 Jan 1;6(1):e003702.
32. African Population and Health Research Center. How Much is Spent to Treat Unsafe Abortion Complications in Kenyan Public Health Facilities?. African Population and Health Research Center; 2018 Apr.

Appendices


Appendix 1: Steps for calculating the induced abortion



$$12 * \left(\frac{\text{typical mo} + \text{last mo}}{2} \right)$$

- PAC cases often undercounted in facility ledgers
- Therefore estimated by asking providers to recall PAC cases seen in a typical month and in the last month
- Remove cases that were initially treated at a facility but then referred to a higher level facility for more care
- Remove late miscarriages (13-24 weeks) which account for 3.41% of live births in reach region

Appendix 2: All pregnancy outcomes in Liberia, 2021



Copyright © 2014 Pearson Education, Inc. All rights reserved. Printed in the United States of America. This publication is protected by copyright. Any unauthorized reproduction or distribution, in any form or by any means, without written permission from Pearson Education, Inc., is prohibited.

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



**African Population and
Health Research Center**

AFRICAN POPULATION AND HEALTH RESEARCH CENTER

Manga Close off Kirawa Road,
P.O. Box 10787-00100,
Nairobi, Kenya

<https://www.aphrc.org/>