

# EVIDENCE BRIEF

## Reducing early and unintended pregnancies among adolescents

→ **Interventions to reduce unmet need for contraception and early and unintended pregnancies among adolescents should be critical components of family planning programmes in developing countries.**

The 1.2 billion adolescents aged 10-19 around the world make up 16% of the world's population (1). The majority (86%) of adolescents live in developing countries. By the time they are 19 years old, half of adolescent girls in developing countries are sexually active, about 40% are married and close to 20% have children (2).

There were 21 million pregnancies among adolescent girls aged 15-19 years in developing countries in 2016; nearly half (49%) were unintended (43% in Asia, 45% in Africa and 74% in Latin America and the Caribbean) (2). An estimated 23 million adolescent girls have an unmet need for modern contraception and are at risk of unintended pregnancy (2). Additionally, 777,000 girls under the age of 15 gave birth in the same year (3).

Early and unintended pregnancy among adolescent girls is influenced by contextual factors at the individual, interpersonal, community and societal levels. It is also associated with adverse health, educational, social and economic outcomes that may impose a substantial burden on the economies and health systems of developing countries (1, 4-10).

Interventions that combine demand-creation activities and provision of contraceptive services have the potential to increase contraceptive uptake among adolescents (11-14). Both the demand for and supply of contraceptives to adolescents can, however, be negatively influenced by several barriers that require appropriate programmatic responses. Table 1 outlines a number of these barriers and responses, and also mentions examples of studies and programmes that used some of these approaches to achieve progress in adolescent uptake of contraception in various regions of the world.



### POLICY AND PROGRAM CONSIDERATIONS

- Collect, analyze and use accurate and up-to-date data.
- Formulate or revise national laws and policies.
- Develop national adolescent sexual and reproductive health strategies.
- Implement strategies with careful monitoring.
- Conduct periodic programme reviews.

**Table 1. Barriers and potential approaches to increase the demand for and supply of contraception among adolescents**

OBJECTIVE	BARRIERS	SUCCESSFUL PROGRAMME APPROACHES	EXAMPLES
<b>DEMAND FOR CONTRACEPTION</b>			
To foster the desire to avoid, delay, space or limit child-bearing	<ul style="list-style-type: none"> <li>Gendered roles (e.g. expectations to be a wife and mother)</li> <li>The need to prove fertility</li> <li>Religious values</li> <li>Norms of the path to adulthood</li> </ul>	Enhance the acceptability of avoiding, delaying, spacing and limiting childbearing.	Conditional cash transfers have transformed life trajectories of girls in Mexico and Malawi.
To foster the desire to use contraception	<ul style="list-style-type: none"> <li>Stigma</li> <li>Taboos (communication and cultural)</li> <li>Lack of understanding (including fear of side-effects)</li> </ul>	Improve the understanding of contraceptive methods and SRH.	Life skills education and vocational training programmes in Uganda and India have increased contraceptive use. In addition, working with influential family members in India helped build support and overcome resistance to adolescent contraceptive use.
To foster a sense of agency in relation to contraceptive use	<ul style="list-style-type: none"> <li>Early marriage</li> <li>Family pressure</li> <li>Sexual coercion and/or violence</li> <li>Limited decision-making autonomy and power</li> </ul>	Increase the sense of agency among girls and women to exert control over their lives and make their own decisions.	Engaging adolescents and their communities directly in Bangladesh and India has improved girls' agency and prevented early marriage.
<b>SUPPLY OF CONTRACEPTION</b>			
To provide access to contraceptive services	<ul style="list-style-type: none"> <li>Lack of awareness of services</li> <li>Inaccessible location</li> <li>Inconvenient operating hours</li> <li>Costs</li> <li>Waiting times</li> </ul>	Increase access to contraceptive services.	Community-based outreach involving provision of information and services through the national Health Extension Program (HEP) led to remarkable improvements in uptake of modern contraception among adolescents in Ethiopia.
To provide adolescent-friendly services	<ul style="list-style-type: none"> <li>Lack of provider sensitivity</li> <li>Provider reluctance to offer contraceptives to adolescents (due to bias)</li> <li>Gender biases</li> <li>Lack of privacy/confidentiality</li> <li>Contraceptives unavailable or out of stock</li> </ul>	Increase provision of high-quality, youth-friendly services for adolescents.	Making services responsive to the needs of adolescents has improved contraceptive use, thereby preventing first pregnancies in China and repeat pregnancies in Kenya. Evidence from studies and projects has been applied at scale in Colombia, Estonia and Malawi.

Source: Adapted from Glinski et al. (2014) (78).

Nine out of ten births to girls aged 15-19 occur within marriage (75). However, in many places sexually active unmarried adolescents have even higher rates of unmet need for contraception (4). The barriers adolescents face in accessing services are often very different for married and unmarried adolescents. Complementary strategies to respond to the differing needs of different populations should be employed in order to leave no one behind.

For example, Estonia implemented a school-based sexuality education programme that linked to youth-friendly sexual and reproductive health (SRH) services and ensured a supportive policy environment. Rates of abortion and births to adolescents aged 15-19 years were substantially reduced (76). Ethiopia achieved remarkable improvements in uptake of modern contraception among adolescents (from less than 10% in 2005 to about 25% in 2011) through the national Health Extension Program (HEP), which involved recruiting, training and deploying an all-female workforce to provide health information and services at the local level (77).

## CONSIDERATIONS FOR REDUCING EARLY AND UNINTENDED PREGNANCIES AMONG ADOLESCENTS

Five elements must be in place in order to apply the evidence to large-scale, national-level programmes:

1. **Collect, analyse and use accurate and up-to-date data** on health outcomes, contraceptive use and its determinants, programme performance and adolescent sexuality/fertility to inform the development of laws, policies and strategies that are responsive to the varying needs of different groups of adolescents based on their social and economic status.
2. **Formulate or revise national laws and policies** to require health workers—in the public, private and non-profit sectors—to provide comprehensive SRH services including contraceptive and safe abortion (where permitted) services to adolescents. Communicate these laws and policies widely.
3. **Develop national adolescent SRH strategies** to include evidence-based and context-specific interventions, budgets to deliver the interventions, and indicators to track progress that are disaggregated by age and socioeconomic status.
4. **Implement strategies with careful monitoring** of activities, and with the input and expertise of civil society groups (including youth organizations and networks).
5. **Conduct periodic programme reviews** to identify lessons learned, build on strengths and address weaknesses.

## REFERENCES

- 1 Adolescents: overview. In: UNICEF data: monitoring the situation of children and women [website]. New York (NY): United Nations Children's Fund; 2016 (<https://data.unicef.org/topic/adolescents/overview/>, accessed 26 June 2017).
- 2 Darroch JE, Woog V, Bankole A, Ashford LS. Adding it up: costs and benefits of meeting contraceptive needs of adolescents. New York (NY): Guttmacher Institute; 2016 ([https://www.guttmacher.org/sites/default/files/report\\_pdf/adding-it-up-adolescents-report.pdf](https://www.guttmacher.org/sites/default/files/report_pdf/adding-it-up-adolescents-report.pdf), accessed 26 June 2017).
- 3 Woog V, Kågesten A. The sexual and reproductive health needs of very young adolescents aged 10-14 in developing countries: what does the evidence show? (2017).
- 4 Woog V, Singh S, Browne A, Philbin J. Adolescent women's need for and use of sexual and reproductive health services in developing countries. New York (NY): Guttmacher Institute; 2015 ([https://www.guttmacher.org/sites/default/files/report\\_pdf/adolescent-srhs-need-developing-countries.pdf](https://www.guttmacher.org/sites/default/files/report_pdf/adolescent-srhs-need-developing-countries.pdf), accessed 26 June 2017).
- 5 Adolescents: health risks and solutions: fact sheet. Geneva: World Health Organization; 2017 (<http://www.who.int/mediacentre/factsheets/fs345/en/>, accessed 26 June 2017).
- 6 Unsafe abortion: global and regional estimates of the incidence of unsafe abortion and associated mortality in 2008, sixth edition. Geneva: World Health Organization; 2011 ([http://www.who.int/reproductivehealth/publications/unsafe\\_abortion/9789241501118/en/](http://www.who.int/reproductivehealth/publications/unsafe_abortion/9789241501118/en/), accessed 26 June 2017).
- 7 Birungi H, Undie C, Mackenzie I, Katahoire A, Obare F, Machawira P. Education sector response to early and unintended pregnancy: a review of country experiences in sub-Saharan Africa. STEP UP and UNESCO Research Report. Strengthening Evidence for Programming on Unintended Pregnancy (STEP UP). Nairobi: Population Council and UNESCO; 2015 ([https://www.popcouncil.org/uploads/pdfs/2015STEPUP\\_EducSectorResp.pdf](https://www.popcouncil.org/uploads/pdfs/2015STEPUP_EducSectorResp.pdf), accessed 26 June 2017).
- 8 Undie C, Birungi H, Odwe G, Obare F. Expanding access to secondary school education for teenage mothers in Kenya: a baseline study report. STEP UP Research Report. Strengthening Evidence for Programming on Unintended Pregnancy (STEP UP). Nairobi: Population Council; 2015 ([https://www.popcouncil.org/uploads/pdfs/2015STEPUP\\_EducTeenMothersKenya.pdf](https://www.popcouncil.org/uploads/pdfs/2015STEPUP_EducTeenMothersKenya.pdf), accessed 26 June 2017).
- 9 Developing an education sector response to early and unintended pregnancy: discussion document for a global consultation. Paris: United Nations Educational, and Scientific and Cultural Organization (UNESCO); 2014 (<http://unesdoc.unesco.org/images/0023/002305/230510E.pdf>, accessed 26 June 2017).
- 10 State of the world population 2013: motherhood in childhood - facing the challenge of adolescent pregnancy. New York (NY): United Nations Population Fund (UNFPA); 2013 (<http://www.unfpa.org/publications/state-world-population-2013>, accessed 26 June 2017).
- 11 Chandra-Mouli V, Lane C, Wong S. What does not work in adolescent sexual and reproductive health: a review of evidence on interventions commonly accepted as best practices. *Glob Health Sci Pract.* 2015; 3(3):333-40. doi:10.9745/GHSP-D-15-00126.

- 12 Gottschalk LB, Ortayli N. Interventions to improve adolescents' contraceptive behaviors in low- and middle-income countries: a review of the evidence base. *Contraception*. 2014; 90(3):211-25. doi:10.1016/j.contraception.2014.04.017.
- 13 Hindin MJ, Kalamar AM, Thompson TA, Upadhyay UD. Interventions to prevent unintended and repeat pregnancy among young people in low- and middle-income countries: a systematic review of the published and gray literature. *J Adol Health*. 2016; 59(suppl. 3):S8-S15. doi:10.1016/j.jadohealth.2016.04.021.
- 14 Youth contraceptive use: effective interventions: a reference guide. Washington (DC): Population Reference Bureau; 2017 (<http://www.prb.org/pdf17/PRB%20Youth%20Policies%20Reference%20Guide.pdf>, accessed 26 June 2017).
- 15 Childhood, not motherhood: preventing adolescent pregnancy. New York: United Nations Population Fund (UNFPA); 2015 ([http://www.unfpa.org/sites/default/files/pub-pdf/Girlhood\\_not\\_motherhood\\_final\\_web.pdf](http://www.unfpa.org/sites/default/files/pub-pdf/Girlhood_not_motherhood_final_web.pdf), accessed 6 July 2017).
- 16 Haldre K, Part K, Ketting E. Youth sexual health improvement in Estonia, 1990–2009: the role of sexuality education and youth-friendly services. *Eur J Contracept Reprod Health Care*. 2012; 17(5):351–62. doi:10.3109/13625187.2012.696751.
- 17 Hounton S, Barros AJD, Amouzou A, Shiferaw A, Maïga A, Akinyemi A et al. Patterns and trends of contraceptive use among sexually active adolescents in Burkina Faso, Ethiopia, and Nigeria: evidence from cross-sectional studies. *Glob Health Action*. 2015; 8:29737. doi:10.3402/gha.v8.29737.
- 18 Glinski A, Sexton M, Petroni S. Understanding the adolescent family planning evidence base. Washington (DC): International Center for Research on Women; 2014.

Authors: Francis Obare (Population Council); Caroline Kabiru, (African Population and Health Research Center); Venkatraman Chandra-Mouli, Matti Parry (World Health Organization).

This is one of seven Family Planning Evidence Briefs prepared for the Family Planning Summit held in London on July 11, 2017. The briefs highlight evidence and provide research and programme considerations for improving access to family planning and reducing unintended pregnancy. Programme considerations are based on the expert views of the authors, who undertook desk reviews drawing on existing evidence.

#### Family Planning Evidence Briefs

- Accelerating uptake of voluntary, rights-based family planning in developing countries (*overview*)
- Ensuring adequate financing of family planning commodities and services
- Reducing early and unintended pregnancies among adolescents
- Improving family planning service delivery in humanitarian crises
- Ensuring contraceptive security through effective supply chains
- Expanding contraceptive choice
- Partnering with the private sector to strengthen provision of contraception

The authors alone are responsible for the views expressed in this article and they do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated.

Family Planning Evidence Brief – Reducing early and unintended pregnancies among adolescents: WHO/RHR/17.10

© World Health Organization 2017. Some rights reserved.  
This work is available under the CC BY-NC-SA 3.0 IGO license.

For more information, please contact: Department of Reproductive Health and Research, World Health Organization, Avenue Appia 20, CH-1211 Geneva 27, Switzerland  
E-mail: [reproductivehealth@who.int](mailto:reproductivehealth@who.int)  
Website: [www.who.int/reproductivehealth](http://www.who.int/reproductivehealth)  
Twitter: @HRPresearch

This material has been funded by UK aid from the UK government; however, the views expressed do not necessarily reflect the UK government's policies.

Prepared July 2017